

## CME

# The Role of the Perioperative Period in Recurrence After Cancer Surgery

Antje Gottschalk, MD,\*† Sonal Sharma, MD,\* Justin Ford, MD,\* Marcel E. Durieux, MD, PhD,\* and Mohamed Tiourine, MD\*

A wealth of basic science data supports the hypothesis that the surgical stress response increases the likelihood of cancer dissemination and metastasis during and after cancer surgery. Anesthetic management of the cancer patient, therefore, could potentially influence long-term outcome. Preclinical data suggest that beneficial approaches might include selection of induction drugs such as propofol, minimizing the use of volatile anesthetics, and coadministration of cyclooxygenase antagonists with systemic opioids. Retrospective clinical trials suggest that the addition of regional anesthesia might decrease recurrence after cancer surgery. Other factors such as blood transfusion, temperature regulation, and statin administration may also affect long-term outcome. (Anesth Analg 2010;110:1636–43)

The idea that surgery promotes local cancer recurrence and distant metastasis is not novel. In fact, >2 millennia ago, observations concerning the negative impact of surgical manipulation on cancer progression were documented. A. Cornelius Celsus, author of “De Medicina,” who established the first staging system of cancer, believed that only *cacoethes* (the first stage of cancer) should be removed, because other stages would be irritated by the treatment.<sup>1</sup> Likewise, Alfred A.L.M. Velpeau (1795–1867) noticed that surgical removal of cancer was associated with the return of the disease and that the operation tended to accelerate tumor growth.<sup>1</sup> Modern therapies such as chemotherapy and radiation have eliminated many cancer fatalities. Nevertheless, despite our advances in cancer treatment technique, metastatic recurrence still remains the leading cause of death from cancer.

Several theories have been advocated to explain the frequent incidence of cancer recurrence, most notably residual minimal disease,<sup>2</sup> dissemination of tumor cells at the time of surgery,<sup>3,4</sup> and possibly tumor dormancy<sup>5</sup> (a period when cancer cells are quiescent before progressive growth). Also, surgery creates profound metabolic, neuroendocrine, inflammatory, and immunological stress.<sup>6–9</sup> This surgical stress response includes the release of chemical mediators that have been directly and indirectly implicated in cancer growth. These mediators could cause an upregulation of major promalignant pathways, resulting in a disruption of

normal tumor homeostasis, thus promoting local and distant metastasis. Importantly, the type of anesthesia may play a role in this process and could indirectly promote malignant cell development. In this article, we briefly review possible mechanisms involved in the effect of surgery on cancer recurrence and then discuss how anesthetic management could potentially influence these mechanisms, thereby affecting long-term patient outcome.

## IMMUNITY AND CANCER

The idea that the immune system recognizes cancerous cells as “nonself” and thereby destroys them was first established by Paul Ehrlich a century ago and was fostered by Burnet and Thomas under the immunosurveillance hypothesis.<sup>10</sup> They suggested that the immune system could actually eliminate cancer cells before they are clinically detectable. It was clear, however, that the immune system is unable to destroy cancer cells completely, as evidenced by the persistence of tumor despite a competent immune system. Subsequently, the concept of immunoediting was born.<sup>10</sup> Under this theory, the immune system is believed to “inadvertently” promote tumor progression by clearing some tumor cells and thereby selecting for those cells most resistant to immune system clearance.

The process of immunoediting is divided into 3 steps. The first step represents the “elimination phase” where cells of the innate and adaptive immune system recognize and destroy tumor cells. The second step is the “equilibrium phase” where it is postulated that the immune system keeps cancer cells in check for a variable period of time. The third step is called “escape” whereby tumor cells evade immunity and become overt tumors.<sup>10</sup> The mechanisms underlying these events are not entirely understood and include alteration in antigen presentation, secretion of immunosuppressive agents, and stimulation of inhibitory pathways. Thus, the immunosuppressant effects of surgery through the secretion of proinflammatory and antiinflammatory cytokines could shape the journey of residual minimal disease toward immune evasion and growth. This hypothesis remains without definitive support at this time.

From the \*Department of Anesthesia, University of Virginia, Charlottesville, Virginia; and †Department of Anesthesiology and Intensive Care Medicine, University Muenster, Muenster, Germany.

Accepted for publication February 24, 2010.

Marcel E. Durieux is section Editor of Anesthetic Pre-Clinical Pharmacology for the Journal. The manuscript was handled by Sorin J. Brull, section Editor of Patient Safety, and Dr. Durieux was not involved in any way with the editorial process or decision.

Disclosure: The authors report no conflict of interest.

Reprints will not be available from the author.

Address correspondence to Mohamed Tiourine, MD, Department of Anesthesiology, University of Virginia, Charlottesville, VA. Address e-mail to mt9y@virginia.edu.

Copyright © 2010 International Anesthesia Research Society  
DOI: 10.1213/ANE.0b013e3181de0ab6

## THE SURGICAL STRESS RESPONSE AND CANCER

Although the primary role of the stress response after surgery is to augment the healing process, either overactivity or underactivity of host-defense mechanisms paradoxically may lead to negative consequences. For instance, the surgical stress response may provide optimal conditions for persistence of residual minimal malignant disease after surgery. Surgery has been suggested to accelerate the development of preexisting micro metastases and to promote the establishment of new metastases.<sup>4,11</sup> It is believed that the postoperative period is the most vulnerable period for potential metastasis after surgery. This vulnerability is mostly attributed to suppression of cell-mediated immunity, the first-line defense mechanism against cancer.<sup>4</sup> The depression of the immune system occurs within hours of surgery, lasts for several days, and is proportional to the extent of surgical trauma.<sup>12</sup> For example, patients with low levels of natural killer (NK) cell activity undergoing primary cancer surgery have been shown to have a higher rate of cancer-related morbidity and mortality. This association has been demonstrated in patients with colorectal,<sup>13,14</sup> gastric,<sup>15</sup> lung,<sup>16</sup> as well as head and neck cancer.<sup>17,18</sup> The underlying mechanisms of postoperative immune suppression have not been completely established; however, existing data strongly suggest a role for the neuroendocrine system, inflammatory system, and the HPA (hypothalamic-pituitary-adrenal) axis.<sup>19</sup>

### Role of the Neuroendocrine System

As early as 1919, scientists demonstrated that young students with pulmonary tuberculosis who were exposed to academic stress had reduced phagocytic capacity to eliminate the pathogen.<sup>20</sup> This was an early demonstration of the effect stress has on the immune system. Likewise, stress has long been considered as a contributor to cancer development.<sup>21</sup> Levels of stress biomarkers, primarily epinephrine and norepinephrine, are elevated in the perioperative period.<sup>19</sup> These neurotransmitters are believed to be responsible for the relationship between stress and cancer progression.<sup>22</sup> This is thought to happen via interaction with  $\beta_1$ - and  $\beta_2$ -adrenergic receptors expressed by tumor cells. Catecholamines have been shown to directly increase the invasive potential of ovarian cancer cells via  $\beta$ -adrenergic upregulation of matrix metalloproteinases<sup>23</sup> and activation of STAT-3 (signal transducer of activation and transcription), a contributor to malignant cell proliferation and survival.<sup>24</sup> Catecholamines also have been shown to increase the production of vascular endothelial growth factor in ovarian cancer cells,<sup>25</sup> and to influence the migration of multiple cancer cell lines including breast, ovarian, and colon cancer.<sup>26</sup> Catecholamines have been shown to influence cell migration and angiogenesis via stimulation of  $\beta_1$  and  $\beta_2$  receptors<sup>27-29</sup> in addition to suppressing cell-mediated immunity (CMI).<sup>4</sup>

### Role of the Inflammatory System

Cancer is often considered an anarchic cell replication process beyond any form of control or recognition; however, this is a profound oversimplification. In fact, cancer growth and metastasis are very complex processes with cell replication merely representing the tip of the iceberg.

Within the tumor microenvironment lies a machinery of highly sophisticated malignant cascades where several products of the inflammatory system such as cytokines, chemokines, prostaglandins (PGs), and cyclooxygenase (COX) are believed to promote cancer progression through immunosuppression, resistance to apoptosis, and promotion of angiogenesis.<sup>30</sup> This is true of the role of chronic inflammation for certain forms of cancer, but it is not clear whether acute inflammation, such as occurs in the perioperative period, results in the same outcomes. Nonetheless, Goldfarb and Ben-Eliyahu<sup>2</sup> suggested that an increase in the level of cytokines (interleukins, IL-6 and IL-8), and PGE<sub>2</sub> in combination with a decrease of T helper type 1 cell-induced cytokine production (IL-2, interferon  $\gamma$ ), could account for the profound suppression of NK cytotoxic activity in the perioperative period.

### Role of the HPA Axis

Pain, a potent stimulant of the HPA axis, has been implicated in causing immunosuppression, making pain management particularly important in the cancer surgery patient. Pain activates the HPA axis and the sympathetic nervous system, thus setting off a cascade of events that leads to immunosuppression. Acute pain has been shown to suppress NK cell activity<sup>31,32</sup> and promote tumor development in animals.<sup>33</sup> Not surprisingly, provision of pain relief has been shown to attenuate surgery-induced increases in metastatic susceptibility in animals.<sup>34</sup>

## POSSIBLE TARGETS FOR METASTASIS PREVENTION BY THE ANESTHESIOLOGIST

Given that the surgical stress response seems to increase opportunities for cancer dissemination and metastasis at the exact time that cancer cells may be released into the circulation, is it possible to minimize these detrimental effects by an appropriate choice of anesthetics? The influence of anesthesia on the stress response to surgery has been investigated in depth. Our goal is to review the anesthetic interventions (Table 1) that affect pathways with a link to tumor progression (Fig. 1).

### Opioids

Opioids have long been the mainstay of treatment of cancer-related pain and are an important modality for the prevention of perioperative pain. As stated above, pain leads to CMI suppression, and its treatment is therefore particularly important. Unfortunately, it also has been established that opioids (morphine in particular) inhibit cellular and humoral immune function in humans.<sup>35-37</sup> There are no data directly implicating opioids in cancer genesis in humans, but animal data strongly suggest that they may contribute to cancer recurrence in the clinical setting. For example, in rodent studies, it was demonstrated that morphine is proangiogenic and promotes breast tumor growth.<sup>38,39</sup> This tumor-promoting effect of opiates was also demonstrated with fentanyl,<sup>40</sup> although in other studies, synthetic opiates did not seem to exhibit immunosuppressive effects. Instead, fentanyl was shown in healthy volunteers to increase NK activity.<sup>41</sup> Morphine has not been demonstrated to be tumor promoting in all models; e.g., it was found to promote apoptosis and cell

**Table 1. Overview of Reported Anesthetic Effects on Tumor Genesis and Recurrence****Surgical stress response and cancer**

Stress and surgical excision of the primary tumor can promote tumor metastasis<sup>4,11</sup>

**Neuroendocrine system**

General anesthesia accompanied by surgical stress may suppress immunity, presumably by directly affecting the immune system or activating the hypothalamic-pituitary-adrenal axis and the sympathetic nervous system<sup>19</sup>

**Inflammatory system**

Promotion of cancer progression through immunosuppression via cytokines, chemokines, prostaglandins, COX<sup>30</sup>

**Pain**

Suppression of NK cell activity<sup>31,32</sup> and promotion of tumor development in animals<sup>33</sup>

**Opioids**

Opioids inhibit cellular and humoral immune function in humans<sup>36</sup>

Morphine inhibits spontaneous and cytokine-enhanced NK cell cytotoxicity<sup>35-37</sup>

In contrast, IV fentanyl increases NK cell cytotoxicity and circulating CD16(+) lymphocytes in humans<sup>41</sup>

Opioid-induced promotion and stimulation of angiogenesis<sup>39</sup>

 **$\beta$ -adrenergic blockade**

$\beta$ -blocker (nadolol) and a prostaglandin synthesis inhibitor (indomethacin) attenuated the metastasis-promoting effects of surgery when used alone or in combination<sup>11</sup>

**COX inhibitors**

COX inhibitors may prevent metastatic progression and attenuate opioid-induced immunosuppression in rats<sup>11</sup>

The combination of COX-2 inhibitor etodolac and  $\beta$ -blocker propranolol can efficiently prevent immunosuppression after surgery<sup>26</sup>

COX-2 inhibitor celecoxib prevents chronic morphine-induced promotion of angiogenesis, tumor growth, metastasis, and mortality in a murine breast cancer model<sup>51</sup>

**Anesthetic induction agents and volatile anesthetics**

Suppression of NK cell activity and promotion of tumor metastasis by ketamine, thiopental, and halothane<sup>63</sup>

**Regional anesthesia**

Studies in animals show that regional anesthesia and optimal postoperative analgesia independently reduce metastasis<sup>8,9</sup>

Retrospective studies in humans support a benefit of regional analgesia for patients undergoing surgery for breast, colon, or prostate cancer with respect to reduction of recurrence<sup>72-74</sup>

**Perioperative blood transfusion**

Perioperative blood transfusion is associated with poorer outcome for patients with colorectal cancer recurrence<sup>82</sup>

**Perioperative hypothermia**

Hypothermia leads to a reduction in cell-mediated immunity, particularly NK cells, and an increase in lung tumor retention and metastasis in rats<sup>87</sup>

COX = cyclooxygenase; NK = natural killer; CD16 = cluster of differentiation.

death in an adenocarcinoma model<sup>42</sup> and in Jurkat cells.<sup>43</sup> Inhibitory effects of morphine on tumor growth have been found in human and animal models as well. For example, in a mouse model, the repeated administration of morphine resulted in decreased tumor cell-induced tissue destruction.<sup>44</sup> This observation was verified in a series of clinical studies showing that pre- and postoperative administration of morphine reduced systemic dissemination of tumor cells.<sup>44-46</sup> This potentially protective effect of morphine on tumor growth was attributed to enhanced T cell-mediated immune responses,<sup>47</sup> physiologically active  $\mu$  opioid receptor splice variant,<sup>48</sup> inhibition of nuclear factor- $\kappa$ B,<sup>49</sup> and nitric oxide release through the constitutive nitric oxide

synthase pathway.<sup>50</sup> Evidently, morphine-tumor interaction is complex, its mechanisms are not completely unraveled, and to a certain extent, contradictory. Further studies in this field are essential to elucidate this interaction. The effects of neuraxial administration of morphine on tumor progression have not been extensively studied. As much smaller doses are used for neuraxial administration, it seems likely that any effects of opioids would be less than after systemic administration.

**COX Inhibitors**

Using a mouse model, Farooqui et al.<sup>51</sup> demonstrated that the chronic use of morphine leads to a stronger expression of COX-2 in tumor cells that increased PG production, which impaired analgesia and increased tumor angiogenesis, growth, metastasis, and mortality.


In addition, they showed that inhibition of COX-2 by administration of celecoxib prevented morphine-induced tumor growth and metastasis and increased survival. This suggests that COX-2 inhibitors may be used in conjunction with opioid to decrease pain in cancer patients, while counterbalancing the negative effects of opioids on immune function. Even in the absence of opiates, COX-2 inhibitors show promise in preventing cancer growth and metastases in animal models. They have multiple effects, including inducing apoptosis,<sup>52-54</sup> decreasing the levels of angiogenic factors,<sup>55-57</sup> and decreasing tumor microvascular density.<sup>26,52</sup>

Indomethacin, a nonselective COX inhibitor, attenuated the metastasis-promoting effect of surgery in rats.<sup>11</sup> Whereas the intraoperative use of indomethacin may be associated with increased blood loss because of COX-1 inhibition, a selective COX-2 inhibitor might be a feasible adjunct that can be used to attenuate perioperative immunosuppression.

In summary, some opiates and COX inhibitors can be used effectively in the cancer surgery patient. The use of celecoxib, a COX-2 inhibitor, is approved by the Food and Drug Administration for the prevention of colorectal cancer in high-risk patients with preexisting susceptibility such as familial adenomatous polyposis. Opioids are probably best used in conjunction with COX inhibitors or when administered via a neuraxial technique. COX inhibitors may prevent metastatic progression and, in addition to their synergistic analgesic properties, attenuate opioid-induced immunosuppression. Although both nonselective COX and selective COX-2 inhibitors are effective, it is recommended that the latter be used to minimize the chance of bleeding and chance of gastric irritation. The risk of cardiovascular complications should additionally be evaluated before their use.

 **$\alpha_2$ -Adrenoceptor Agonists**

Since the 1980s, clonidine, an antihypertensive drug with sedative properties, has been used as an adjuvant to local anesthetics in various regional techniques to extend the duration of the block.<sup>58</sup> Dexmedetomidine is indicated for sedation in patients receiving mechanical ventilation in the intensive care unit. In 2006, Vázquez et al.<sup>59</sup> published the first study to describe the presence of  $\alpha_2$ -adrenoceptors in human epithelial breast cell lines. In 2008, Bruzzone et al.<sup>60</sup> demonstrated a significant enhancement of mouse mammary tumor growth induced by clonidine. In that study, incubation for 2 days with the  $\alpha_2$ -adrenoceptor agonists



**Figure 1.** Perioperative mechanism of immunosuppression. STAT = signal transducer of activation and transcription; COX = cyclooxygenase; NK = natural killer.

(clonidine  $0.1 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{d}^{-1}$  and dexmedetomidine  $0.05 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{d}^{-1}$ ) significantly enhanced proliferation of the mammary tumor cells. The  $\alpha_2$ -adrenoceptor antagonists yohimbine ( $0.5 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{d}^{-1}$ ) and rauwolscine ( $0.5 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{d}^{-1}$ ) completely reversed the enhancement of tumor growth of clonidine. These results suggest that the  $\alpha_2$ -agonist clonidine exerts its enhancement on tumor growth by enhancing cell proliferation and decreasing cell apoptosis, whereas the inverse agonist (i.e., an agent that exerts the opposite pharmacologic effect of an agonist) rauwolscine exerts its protective action both by enhancing apoptosis and reducing cell proliferation. However, the group receiving yohimbine alone showed a nonsignificant but constant increase in tumor growth, whereas rauwolscine alone diminished tumor growth significantly, behaving as a reverse agonist. Therefore, the possibility of blocking this  $\alpha_2$ -adrenoceptor-mediated tumor enhancement by hormones released during stress could be an interesting adjuvant therapy for breast cancer patients.<sup>60</sup>

### $\beta$ -Adrenergic Blockade

Hasegawa and Saiki<sup>61</sup> demonstrated a relationship among stress, tumor growth, and  $\beta$ -adrenergic activation independent of glucocorticoid levels in mice. They also showed that the administration of  $\beta$ -blockers abrogated this effect. A combination of  $\beta$ -blockers and COX-2 inhibitors improves immune competence and reduces the risk of tumor metastasis after surgery in animals.<sup>26</sup> Palm et al.<sup>62</sup> demonstrated that use of  $\beta$ -blockers in mice with prostate carcinoma inhibited lumbar lymph node metastases. It seems that  $\beta$ -blockers act by inhibition of catecholamine's induced

signal transducer and activator of transcription 3 (STAT-3) activity. STATs are a family of transcription factors that regulate the expression of certain immune system genes. Given the available animal data on the effect of catecholamine on cancer progression and the chemopreventive effect of  $\beta$ -blockers, it seems possible that  $\beta$ -blockers may be beneficial in preventing metastatic progression in humans.

### Anesthetic Induction Drugs and Volatile Anesthetics

Melamed et al.<sup>63</sup> demonstrated in rats that ketamine, thiopental, and halothane reduced NK cell activity and increased lung tumor retention and metastasis. The number of circulating NK cells per milliliter of blood was reduced significantly by ketamine and thiopental. The effect of halothane was similar, but did not reach statistical significance.<sup>63</sup> The authors suggested that a reduction in NK cell activity might have a major impact on the resistance to tumor development. The interaction of ketamine with  $\alpha$ - and  $\beta$ -adrenoreceptors could be one reason for the suppression of NK activity and the promotion of metastasis.<sup>64</sup> Although the mechanism for immune suppression induced by thiopental or volatile anesthetics remains to be elucidated, it is not related to the anesthetic state per se.<sup>63</sup> Of interest, propofol did not exhibit these effects; instead, propofol seems to have protective effects through various mechanisms, including inhibition of COX-2,<sup>65</sup> inhibition of PGE<sub>2</sub>, but also through enhancement of antitumor immunity.<sup>66</sup> Ke et al.<sup>67</sup> showed in patients undergoing open cholecystectomy that the combination of propofol and remifentanyl

resulted in an increase in antiinflammatory cytokines IL-10 (known to have antitumor activity and help with healing and repair), as compared with inhaled anesthesia with isoflurane. Inada et al.<sup>68</sup> found in patients undergoing supratentorial tumor excision that propofol anesthesia mitigated the adverse effects of surgical stress-induced immune response better than isoflurane. It would be intuitive therefore to think that total IV anesthesia is preferable to an inhaled technique for patients undergoing cancer surgery.

Although less information is available on other drugs used for anesthesia induction, they may affect the immune system as well. For example, midazolam decreases IL-8 levels. This may contribute to immunosuppression because IL-8 is a chemotactic and activating factor that mediates neutrophil adhesion and margination and is essential for host defense.<sup>69</sup>

### Regional Anesthesia

Regional anesthesia, including spinal and epidural anesthesia, reduces the stress response caused by surgery, which is believed to be a mediator of postoperative immunosuppression.<sup>70,71</sup> Regional anesthesia attenuates the surgical stress response by blocking afferent neural transmission. This prevents noxious afferent input from reaching the central nervous system. The addition of regional anesthesia to general anesthesia also results in less overall use of opioids and volatile anesthetics. This association may be beneficial to patients undergoing cancer surgery, because it should theoretically result in less immunosuppression.

In a retrospective analysis of patients undergoing surgical treatment for breast cancer, Exadaktylos et al.<sup>72</sup> demonstrated that the use of paravertebral nerve block in combination with general anesthesia was associated with a longer cancer-free interval and a lower incidence of recurrence. Similar results were shown in other retrospective studies of the use of epidural local anesthetics in prostate cancer and colon cancer surgery.<sup>73,74</sup> Moreover, in a large-scale study in patients with melanoma, substitution of local anesthesia for general anesthesia independently predicted a decrease in tumor recurrence.<sup>75</sup> Also, a recent study of the effect of paravertebral blocks and propofol on cytokine response during breast cancer surgery noted a decrease in tumorigenic cytokines IL-1 $\beta$ /IL-8 and an increase in IL-10, a known antitumor cytokine.<sup>76</sup> The results of these studies should be interpreted cautiously, and there still is no clear evidence whether a simple change in anesthetic practice could affect patient survival. Several multicenter prospective randomized controlled trials are underway, which will test the hypothesis that local or metastatic recurrence after several types of cancer surgery will be decreased in patients randomized to a regional anesthetic technique in comparison to those receiving general anesthesia.<sup>77</sup>

### Anemia and Perioperative Blood Transfusion

Anemia is an ominous sign for the cancer patient. Caro et al.<sup>78</sup> conducted a systematic quantitative review and demonstrated that anemia is associated with increased postoperative morbidity and mortality in all forms of cancer, with decreased local control in surgically treated squamous cell carcinoma<sup>79</sup> and decreased survival in non-small cell lung carcinoma.<sup>80</sup> It may seem logical that transfusing the anemic cancer patient would lead to an increase in survival;

however, evidence suggests that transfusion poses an independent risk to the cancer patient.

In 1973, Opelz et al.<sup>81</sup> proposed the idea of transfusion-related immunomodulation (TRIM) after recognizing that kidney transplant recipients who received >10 U of allogenic blood had better allograft survival. The demonstration of TRIM has led many to hypothesize that patients undergoing surgery for cancer are at an increased risk for metastatic recurrence if they receive blood products. A recent meta-analysis conducted by Amato and Pescatori<sup>82</sup> suggests this may be correct for patients with colorectal cancer. The authors reviewed 36 studies involving >12,000 patients and found a moderate association between perioperative blood transfusion and colorectal cancer recurrence with an odds ratio of 1.42 (95% confidence interval, 1.20–1.67). These findings call for carefully restricted indications for perioperative transfusions in colorectal cancer patients operated on for cure, and await the results of future studies addressing the role of surgery-related risk factors on the need for transfusion and disease recurrence.<sup>82</sup> However, perioperative transfusion requirement and transfusion within 30 days of operation were not significant predictors of survival for patients with pancreatic ductal adenocarcinoma.<sup>83</sup>

There are >200 published reports on transfusion-related metastatic recurrence with no official consensus. In fact, the only consistency among published data is that cancer patients who receive blood transfusions during surgery tend to do worse. This is independent of whether the blood is allogenic, autogenic, or leukocyte reduced. We are still awaiting human data with regard to the age of the blood products; however, a link between cancer progression and aged erythrocytes was recently demonstrated in a rat model.<sup>84</sup> Nevertheless, it seems that both anemia and blood transfusions are associated with poorer outcomes in cancer patients. Perhaps factors influencing the need for blood transfusion have a greater bearing on prognosis than the receipt of blood itself. For the anesthesiologist, it is important to ensure that the patient is medically optimized before surgery, that all attempts are made to perform minimal blood loss surgery, and that transfusions are used judiciously.

### Perioperative Hypothermia

Perioperative hypothermia is associated with an increase in wound infections.<sup>85</sup> In fact, maintaining normothermia is more effective than perioperative antibiotics in the prevention of wound infections.<sup>85</sup> Hypothermia also causes increased blood loss and predisposes patients to blood transfusions.<sup>86</sup> Additionally, hypothermia in combination with surgery and general anesthesia has been shown to lead to a reduction in CMI, particularly NK cells, and an increase in lung tumor retention and metastasis in rats.<sup>87</sup> Although a retrospective analysis could not confirm this finding in humans,<sup>88</sup> it seems important that the anesthesia provider maintains normothermia in cancer patients. This simple measure may lead to a reduction in cancer recurrence after surgery and will most assuredly lead to a reduction in perioperative infections, blood loss, and need for blood transfusion.

## Statins

Statins first attracted interest for cancer prevention as an unexpected result of safety monitoring in large randomized controlled trials of statins' and other lipid-decreasing drugs' effectiveness in preventing cardiovascular disease. This monitoring was implemented because the randomized controlled trials showed consistent increases in statin-associated noncardiovascular disease mortality.<sup>89</sup> The safety results, however, indicated that statins did not increase cancer incidence or cancer mortality. In fact, subsequent intensive clinical and observational studies and preclinical data showed significant statin-associated reduction in overall cancer incidence, the most promising of which are colorectal, prostate, breast, and skin cancers.<sup>90,91</sup> This has been further highlighted by 2 large population-based studies that showed statin-associated reduction in the risk of colorectal and advanced prostate cancers.<sup>92</sup>

The effects of statins have been shown to occur through 3-hydroxy-3-methylglutaryl coenzyme A reductase-dependent or -independent pathways such as binding to lymphocyte function associated antigen 1, which has an important role in leukocyte migration and T cell activation.<sup>93</sup> One study of ischemia and cancer indicated that statins augmented blood flow to the ischemic tissue but did not increase blood flow or capillary density in implanted colon tumors. In fact, the growth of these blood vessels was substantially delayed, suggesting an antiangiogenic effect of statins in carcinogenesis.<sup>94</sup> The beneficial effects of statins on carcinogenesis have also been linked to their antiinflammatory and immunomodulatory effects on adhesion, inflammatory mediators, major histocompatibility complex II, T helper 1 and 2 cytokines, and C-reactive protein.<sup>95</sup> Statins also seem to induce apoptosis and inhibit proliferation by regulating several signaling pathways in malignant cells.<sup>96</sup> This pleiotropic aspect of statins indicates the broad impact that these drugs can have on public health, which should be defined by well-designed observational studies of cancer within large prospective cohorts.

## CONCLUSIONS

It is much too early to write evidence-based guidelines for the anesthesiologist's role in the prevention of cancer recurrence. At this time, we have very interesting animal data (but do not know if these can be extrapolated to medical practice) and very limited, mostly retrospective, clinical studies. Many questions remain unanswered: How much anesthetic exposure is necessary? Do the type, grade, stage, and location of tumor matter? How can we optimally influence the inflammatory response in the postoperative period? As anesthesiologists, we render care to patients coming to the operating room for diverse and complex cancer operations. By this time, the tumor has already grown, micrometastases already exist, tumor manipulation will cause malignant cells to migrate, and in many patients recurrence will occur after a variable period of time. The possibility that perioperative management may alter the rate or incidence of recurrence is tremendously exciting, but much more research is needed in order for this possibility to be conclusively demonstrated. ■■

## REFERENCES

1. Raven R. *The Theory and Practice of Oncology: Historical Evolution and Present Principles*. Park Ridge, NJ: Parthenon Publishing Group, 1990
2. Goldfarb Y, Ben-Eliyahu S. Surgery as a risk factor for breast cancer recurrence and metastasis: mediating mechanisms and clinical prophylactic approaches. *Breast Dis* 2006;26:99–114
3. Yamashita JI, Kurusu Y, Fujino N, Saisyoji T, Ogawa M. Detection of circulating tumor cells in patients with non-small cell lung cancer undergoing lobectomy by video-assisted thoracic surgery: a potential hazard for intraoperative hematogenous tumor cell dissemination. *J Thorac Cardiovasc Surg* 2000;119:899–905
4. Ben-Eliyahu S. The promotion of tumor metastasis by surgery and stress: immunological basis and implications for psychoneuroimmunology. *Brain Behav Immun* 2003;17(Suppl 1):S27–36
5. Demicheli R, Miceli R, Moliterni A, Zambetti M, Hrushesky WJ, Retsky MW, Valagussa P, Bonadonna G. Breast cancer recurrence dynamics following adjuvant CMF is consistent with tumor dormancy and mastectomy-driven acceleration of the metastatic process. *Ann Oncol* 2005;16:1449–57
6. Rosendahl W, Schulz U, Teufel T, Irtel von Brenndorf C, Gupta D. Surgical stress and neuroendocrine responses in infants and children. *J Pediatr Endocrinol Metab* 1995;8:187–94
7. Kennedy BC, Hall GM. Neuroendocrine and inflammatory aspects of surgery: do they affect outcome? *Acta Anaesthesiol Belg* 1999;50:205–9
8. Wada H, Seki S, Takahashi T, Kawarabayashi N, Higuchi H, Habu Y, Sugahara S, Kazama T. Combined spinal and general anesthesia attenuates liver metastasis by preserving TH1/TH2 cytokine balance. *Anesthesiology* 2007;106:499–506
9. Bar-Yosef S, Melamed R, Page GG, Shakhar K, Shakhar K, Ben-Eliyahu S. Attenuation of the tumor-promoting effect of surgery by spinal blockade in rats. *Anesthesiology* 2001;94:1066–73
10. Dunn GP, Old LJ, Schreiber RD. The immunobiology of cancer immunosurveillance and immunoediting. *Immunity* 2004;21:137–48
11. Melamed R, Rosenne E, Shakhar K, Schwartz Y, Abudarham N, Ben-Eliyahu S. Marginating pulmonary-NK activity and resistance to experimental tumor metastasis: suppression by surgery and the prophylactic use of a beta-adrenergic antagonist and a prostaglandin synthesis inhibitor. *Brain Behav Immun* 2005;19:114–26
12. Page GG. Surgery-induced immunosuppression and postoperative pain management. *AACN Clin Issues* 2005;16:302–9
13. Tartter PJ, Steinberg B, Barron DM, Martinelli G. The prognostic significance of natural killer cytotoxicity in patients with colorectal cancer. *Arch Surg* 1987;122:1264–8
14. Koda K, Saito N, Takiguchi N, Oda K, Nunomura M, Nakajima N. Preoperative natural killer cell activity: correlation with distant metastases in curatively resected colorectal carcinomas. *Int Surg* 1997;82:190–3
15. Takeuchi H, Maehara Y, Tokunaga E, Koga T, Kakeji Y, Sugimachi K. Prognostic significance of natural killer cell activity in patients with gastric carcinoma: a multivariate analysis. *Am J Gastroenterol* 2001;96:574–8
16. Fujisawa T, Yamaguchi Y. Autologous tumor killing activity as a prognostic factor in primary resected non-small cell carcinoma of the lung. *Cancer* 1997;79:474–81
17. Brittenden J, Heys SD, Ross J, Eremin O. Natural killer cells and cancer. *Cancer* 1996;77:1226–43
18. Schantz SP, Savage HE, Racz T, Liu FJ, Brown BW, Rossen RD, Hong WK. Immunologic determinants of head and neck cancer response to induction chemotherapy. *J Clin Oncol* 1989;7:857–64
19. Kurosawa S, Kato M. Anesthetics, immune cells, and immune responses. *J Anesth* 2008;22:263–77
20. Ishigami T. The influence of psychic acts on the progress of pulmonary tuberculosis. *Am Rev Tuberc* 1919;2:470–84
21. Reiche EM, Nunes SO, Morimoto HK. Stress, depression, the immune system, and cancer. *Lancet Oncol* 2004;5:617–25
22. Thaker PH, Sood AK. Neuroendocrine influences on cancer biology. *Semin Cancer Biol* 2008;18:164–70

23. Sood AK, Bhatti R, Kamat AA, Landen CN, Han L, Thaker PH, Li Y, Gershenson DM, Lutgendorf S, Cole SW. Stress hormone-mediated invasion of ovarian cancer cells. *Clin Cancer Res* 2006;12:369–75
24. Landen CN Jr, Lin YG, Armaiz Pena GN, Das PD, Arevalo JM, Kamat AA, Han LY, Jennings NB, Spannuth WA, Thaker PH, Lutgendorf SK, Savary CA, Sanguino AM, Lopez-Berestein G, Cole SW, Sood AK. Neuroendocrine modulation of signal transducer and activator of transcription-3 in ovarian cancer. *Cancer Res* 2007;67:10389–96
25. Lutgendorf SK, Cole S, Costanzo E, Bradley S, Coffin J, Jabbari S, Rainwater K, Ritchie JM, Yang M, Sood AK. Stress-related mediators stimulate vascular endothelial growth factor secretion by two ovarian cancer cell lines. *Clin Cancer Res* 2003;9:4514–21
26. Benish M, Bartal I, Goldfarb Y, Levi B, Avraham R, Raz A, Ben-Eliyahu S. Perioperative use of beta-blockers and COX-2 inhibitors may improve immune competence and reduce the risk of tumor metastasis. *Ann Surg Oncol* 2008;15:2042–52
27. Yang EV, Sood AK, Chen M, Li Y, Eubank TD, Marsh CB, Jewell S, Flavahan NA, Morrison C, Yeh PE, Lemeshow S, Glaser R. Norepinephrine up-regulates the expression of vascular endothelial growth factor, matrix metalloproteinase (MMP)-2, and MMP-9 in nasopharyngeal carcinoma tumor cells. *Cancer Res* 2006;66:10357–64
28. Thaker PH, Han LY, Kamat AA, Arevalo JM, Takahashi R, Lu C, Jennings NB, Armaiz-Pena G, Bankson JA, Ravoori M, Merritt WM, Lin YG, Mangala LS, Kim TJ, Coleman RL, Landen CN, Li Y, Felix E, Sanguino AM, Newman RA, Lloyd M, Gershenson DM, Kundra V, Lopez-Berestein G, Lutgendorf SK, Cole SW, Sood AK. Chronic stress promotes tumor growth and angiogenesis in a mouse model of ovarian carcinoma. *Nat Med* 2006;12:939–44
29. Masur K, Niggemann B, Zanker KS, Entschladen F. Norepinephrine-induced migration of SW 480 colon carcinoma cells is inhibited by beta-blockers. *Cancer Res* 2001;61:2866–9
30. Kundu JK, Surh YJ. Inflammation: gearing the journey to cancer. *Mutat Res* 2008;659:15–30
31. Sacerdote P, Manfredi B, Bianchi M, Panerai AE. Intermittent but not continuous inescapable footshock stress affects immune responses and immunocyte beta-endorphin concentrations in the rat. *Brain Behav Immun* 1994;8:251–60
32. Shavit Y, Martin FC, Yirmiya R, Ben-Eliyahu S, Terman GW, Weiner H, Gale RP, Liebeskind JC. Effects of a single administration of morphine or footshock stress on natural killer cell cytotoxicity. *Brain Behav Immun* 1987;1:318–28
33. Lewis JW, Shavit Y, Terman GW, Nelson LR, Gale RP, Liebeskind JC. Apparent involvement of opioid peptides in stress-induced enhancement of tumor growth. *Peptides* 1983;4:635–8
34. Page GG, Blakely WP, Ben-Eliyahu S. Evidence that postoperative pain is a mediator of the tumor-promoting effects of surgery in rats. *Pain* 2001;90:191–9
35. Yeager MP, Colacchio TA, Yu CT, Hildebrandt L, Howell AL, Weiss J, Guyre PM. Morphine inhibits spontaneous and cytokine-enhanced natural killer cell cytotoxicity in volunteers. *Anesthesiology* 1995;83:500–8
36. Sacerdote P, Bianchi M, Gaspani L, Manfredi B, Maucione A, Terno G, Ammatuna M, Panerai AE. The effects of tramadol and morphine on immune responses and pain after surgery in cancer patients. *Anesth Analg* 2000;90:1411–4
37. Beilin B, Shavit Y, Hart J, Mordashov B, Cohn S, Notti I, Bessler H. Effects of anesthesia based on large versus small doses of fentanyl on natural killer cell cytotoxicity in the perioperative period. *Anesth Analg* 1996;82:492–7
38. Ben-Eliyahu S, Page GG, Yirmiya R, Shakhar G. Evidence that stress and surgical interventions promote tumor development by suppressing natural killer cell activity. *Int J Cancer* 1999;80:880–8
39. Gupta K, Kshirsagar S, Chang L, Schwartz R, Law P, Yee D, Heibel R. Morphine stimulates angiogenesis by activating proangiogenic and survival-promoting signaling and promotes breast tumor growth. *Cancer Res* 2002;62:4491–8
40. Shavit Y, Ben-Eliyahu S, Zeidel A, Beilin B. Effects of fentanyl on natural killer cell activity and on resistance to tumor metastasis in rats. Dose and timing study. *Neuroimmunomodulation* 2004;11:255–60
41. Yeager MP, Procopio MA, DeLeo JA, Arruda JL, Hildebrandt L, Howell AL. Intravenous fentanyl increases natural killer cell cytotoxicity and circulating CD16(+) lymphocytes in humans. *Anesth Analg* 2002;94:94–9
42. Tegeger I, Grosch S, Schmidtko A, Haussler A, Schmidt H, Niederberger E, Scholich K, Geisslinger G. G protein-independent G1 cell cycle block and apoptosis with morphine in adenocarcinoma cells: involvement of p53 phosphorylation. *Cancer Res* 2003;63:1846–52
43. Singhal PC, Kapasi AA, Reddy K, Franki N, Gibbons N, Ding G. Morphine promotes apoptosis in Jurkat cells. *J Leukoc Biol* 1999;66:650–8
44. Page GG, Ben-Eliyahu S, Yirmiya R, Liebeskind JC. Morphine attenuates surgery-induced enhancement of metastatic colonization in rats. *Pain* 1993;54:21–8
45. Page GG, Ben-Eliyahu S, Liebeskind JC. The role of LGL/NK cells in surgery-induced promotion of metastasis and its attenuation by morphine. *Brain Behav Immun* 1994;8:241–50
46. Page GG, McDonald JS, Ben-Eliyahu S. Pre-operative versus postoperative administration of morphine: impact on the neuroendocrine, behavioural, and metastatic-enhancing effects of surgery. *Br J Anaesth* 1998;81:216–23
47. Fuggetta MP, Di Francesco P, Falchetti R, Cottarelli A, Rossi L, Tricarico M, Lanzilli G. Effect of morphine on cell-mediated immune responses of human lymphocytes against allogeneic malignant cells. *J Exp Clin Cancer Res* 2005;24:255–63
48. Cadet P, Mantione KJ, Stefano GB. Molecular identification and functional expression of mu 3, a novel alternatively spliced variant of the human mu opiate receptor gene. *J Immunol* 2003;170:5118–23
49. Sueoka E, Sueoka N, Kai Y, Okabe S, Suganuma M, Kanematsu K, Yamamoto T, Fujiki H. Anticancer activity of morphine and its synthetic derivative, KT-90, mediated through apoptosis and inhibition of NF-kappaB activation. *Biochem Biophys Res Commun* 1998;252:566–70
50. Welters ID, Menzebach A, Goumon Y, Cadet P, Menges T, Hughes TK, Hempelmann G, Stefano GB. Morphine inhibits NF-kappaB nuclear binding in human neutrophils and monocytes by a nitric oxide-dependent mechanism. *Anesthesiology* 2000;92:1677–84
51. Farooqui M, Li Y, Rogers T, Poonawala T, Griffin RJ, Song CW, Gupta K. COX-2 inhibitor celecoxib prevents chronic morphine-induced promotion of angiogenesis, tumour growth, metastasis and mortality, without compromising analgesia. *Br J Cancer* 2007;97:1523–31
52. Roche-Nagle G, Connolly EM, Eng M, Bouchier-Hayes DJ, Harmey JH. Antimetastatic activity of a cyclooxygenase-2 inhibitor. *Br J Cancer* 2004;91:359–65
53. Sinicropo FA, Gill S. Role of cyclooxygenase-2 in colorectal cancer. *Cancer Metastasis Rev* 2004;23:63–75
54. Kern MA, Haugg AM, Koch AF, Schilling T, Breuhahn K, Walczak H, Fleischer B, Trautwein C, Michalski C, Schulze-Bergkamen H, Friess H, Stremmel W, Krammer PH, Schirmacher P, Muller M. Cyclooxygenase-2 inhibition induces apoptosis signaling via death receptors and mitochondria in hepatocellular carcinoma. *Cancer Res* 2006;66:7059–66
55. Wei D, Wang L, He Y, Xiong HQ, Abbruzzese JL, Xie K. Celecoxib inhibits vascular endothelial growth factor expression in and reduces angiogenesis and metastasis of human pancreatic cancer via suppression of Sp1 transcription factor activity. *Cancer Res* 2004;64:2030–8
56. Jones MK, Wang H, Peskar BM, Levin E, Itani RM, Sarfeh II, Tarnawski AS. Inhibition of angiogenesis by nonsteroidal anti-inflammatory drugs: insight into mechanisms and implications for cancer growth and ulcer healing. *Nat Med* 1999;5:1418–23
57. Rozic JG, Chakraborty C, Lala PK. Cyclooxygenase inhibitors retard murine mammary tumor progression by reducing tumor cell migration, invasiveness and angiogenesis. *Int J Cancer* 2001;93:497–506
58. Duma A, Urbanek B, Sitzwohl C, Kreiger A, Zimpfer M, Kapral S. Clonidine as an adjuvant to local anaesthetic axillary brachial plexus block: a randomized, controlled study. *Br J Anaesth* 2005;94:112–6

59. Vázquez SM, Mladovan AG, Pérez C, Bruzzone A, Baldi A, Lüthy IA. Human breast cell lines exhibit functional alpha2-adrenoceptors. *Cancer Chemother Pharmacol* 2006;58:50–61
60. Bruzzone A, Pinero CP, Castillo LF, Sarappa MG, Rojas P, Lanari C, Luthy IA. Alpha2-adrenoceptor action on cell proliferation and mammary tumour growth in mice. *Br J Pharmacol* 2008;155:494–504
61. Hasegawa H, Saiki I. Psychosocial stress augments tumor development through beta-adrenergic activation in mice. *Jpn J Cancer Res* 2002;93:729–35
62. Palm D, Lang K, Niggemann B, Drell TL IV, Masur K, Zaenker KS, Entschladen F. The norepinephrine-driven metastasis development of PC-3 human prostate cancer cells in BALB/c nude mice is inhibited by beta-blockers. *Int J Cancer* 2006;118:2744–9
63. Melamed R, Bar-Yosef S, Shakhar G, Shakhar K, Ben-Eliyahu S. Suppression of natural killer cell activity and promotion of tumor metastasis by ketamine, thiopental, and halothane, but not by propofol: mediating mechanisms and prophylactic measures. *Anesth Analg* 2003;97:1331–9
64. Shakhar G, Ben-Eliyahu S. In vivo beta-adrenergic stimulation suppresses natural killer cell activity and compromises resistance to tumor metastasis in rats. *J Immunol* 1998;160:3251–8
65. Inada T, Kubo K, Kambara T, Shingu K. Propofol inhibits cyclo-oxygenase activity in human monocytic THP-1 cells. *Can J Anaesth* 2009;56:222–9
66. Kushida A, Inada T, Shingu K. Enhancement of antitumor immunity after propofol treatment in mice. *Immunopharmacol Immunotoxicol* 2007;29:477–86
67. Ke JJ, Zhan J, Feng XB, Wu Y, Rao Y, Wang YL. A comparison of the effect of total intravenous anaesthesia with propofol and remifentanyl and inhalational anaesthesia with isoflurane on the release of pro- and anti-inflammatory cytokines in patients undergoing open cholecystectomy. *Anaesth Intensive Care* 2008;36:74–8
68. Inada T, Yamanouchi Y, Jomura S, Sakamoto S, Takahashi M, Kambara T, Shingu K. Effect of propofol and isoflurane anaesthesia on the immune response to surgery. *Anaesthesia* 2004;59:954–9
69. Galley HF, Dubbels AM, Webster NR. The effect of midazolam and propofol on interleukin-8 from human polymorphonuclear leukocytes. *Anesth Analg* 1998;86:1289–93
70. Liu SS, Block BM, Wu CL. Effects of perioperative central neuraxial analgesia on outcome after coronary artery bypass surgery: a meta-analysis. *Anesthesiology* 2004;101:153–61
71. Tonnesen E, Wahlgreen C. Influence of extradural and general anaesthesia on natural killer cell activity and lymphocyte subpopulations in patients undergoing hysterectomy. *Br J Anaesth* 1988;60:500–7
72. Exadaktylos AK, Buggy DJ, Moriarty DC, Mascha E, Sessler DI. Can anesthetic technique for primary breast cancer surgery affect recurrence or metastasis? *Anesthesiology* 2006;105:660–4
73. Biki B, Mascha E, Moriarty DC, Fitzpatrick JM, Sessler DI, Buggy DJ. Anesthetic technique for radical prostatectomy surgery affects cancer recurrence: a retrospective analysis. *Anesthesiology* 2008;109:180–7
74. Christopherson R, James KE, Tableman M, Marshall P, Johnson FE. Long-term survival after colon cancer surgery: a variation associated with choice of anesthesia. *Anesth Analg* 2008;107:325–32
75. Schlagenhauff B, Ellwanger U, Breuninger H, Stroebel W, Rassner G, Garbe C. Prognostic impact of the type of anaesthesia used during the excision of primary cutaneous melanoma. *Melanoma Res* 2000;10:165–9
76. Deegan CA, Buggy DJ, Moriarty DC, Sessler DI, Kavanagh DP. Anesthetic technique and the cytokine response to primary breast cancer surgery. *ASA annual meeting, Orlando, FL, 2008:A841*
77. Sessler DI, Ben-Eliyahu S, Mascha EJ, Parat MO, Buggy DJ. Can regional analgesia reduce the risk of recurrence after breast cancer? Methodology of a multicenter randomized trial. *Contemp Clin Trials* 2008;29:517–26
78. Caro JJ, Salas M, Ward A, Goss G. Anemia as an independent prognostic factor for survival in patients with cancer: a systematic, quantitative review. *Cancer* 2001;91:2214–21
79. Lutterbach J, Guttenberger R. Anemia is associated with decreased local control of surgically treated squamous cell carcinomas of the glottic larynx. *Int J Radiat Oncol Biol Phys* 2000;48:1345–50
80. Yovino S, Kwok Y, Krasna M, Bangalore M, Suntharalingam M. An association between preoperative anemia and decreased survival in early-stage non-small-cell lung cancer patients treated with surgery alone. *Int J Radiat Oncol Biol Phys* 2005;62:1438–43
81. Opelz G, Sengar DP, Mickey MR, Terasaki PI. Effect of blood transfusions on subsequent kidney transplants. *Transplant Proc* 1973;5:253–9
82. Amato A, Pescatori M. Perioperative blood transfusions for the recurrence of colorectal cancer. *Cochrane Database Syst Rev* 2006;1:CD005033
83. Clark E, Connor S, Taylor MA, Hendry CL, Madhavan KK, Garden OJ, Parks RW. Perioperative transfusion for pancreaticoduodenectomy and its impact on prognosis in resected pancreatic ductal adenocarcinoma. *HPB (Oxford)* 2007;9:472–7
84. Atzil S, Arad M, Glasner A, Abiri N, Avraham R, Greenfeld K, Rosenne E, Beilin B, Ben-Eliyahu S. Blood transfusion promotes cancer progression: a critical role for aged erythrocytes. *Anesthesiology* 2008;109:989–97
85. Kurz A, Sessler DI, Lenhardt R. Perioperative normothermia to reduce the incidence of surgical-wound infection and shorten hospitalization. Study of Wound Infection and Temperature Group. *N Engl J Med* 1996;334:1209–15
86. Rajagopalan S, Mascha E, Na J, Sessler DI. The effects of mild perioperative hypothermia on blood loss and transfusion requirement. *Anesthesiology* 2008;108:71–7
87. Ben-Eliyahu S, Shakhar G, Rosenne E, Levinson Y, Beilin B. Hypothermia in barbiturate-anesthetized rats suppresses natural killer cell activity and compromises resistance to tumor metastasis: a role for adrenergic mechanisms. *Anesthesiology* 1999;91:732–40
88. Yucel Y, Barlan M, Lenhardt R, Kurz A, Sessler DI. Perioperative hypothermia does not enhance the risk of cancer dissemination. *Am J Surg* 2005;189:651–5
89. Oliver MF. Might treatment of hypercholesterolaemia increase non-cardiac mortality? *Lancet* 1991;337:1529–31
90. Graaf MR, Richel DJ, van Noorden CJ, Guchelaar HJ. Effects of statins and farnesyltransferase inhibitors on the development and progression of cancer. *Cancer Treat Rev* 2004;30:609–41
91. Blais L, Desgagne A, LeLorier J. 3-Hydroxy-3-methylglutaryl coenzyme A reductase inhibitors and the risk of cancer: a nested case-control study. *Arch Intern Med* 2000;160:2363–8
92. Poynter JN, Gruber SB, Higgins PD, Almog R, Bonner JD, Rennert HS, Low M, Greenson JK, Rennert G. Statins and the risk of colorectal cancer. *N Engl J Med* 2005;352:2184–92
93. Weitz-Schmidt G, Welzenbach K, Brinkmann V, Kamata T, Kallen J, Bruns C, Cottens S, Takada Y, Hommel U. Statins selectively inhibit leukocyte function antigen-1 by binding to a novel regulatory integrin site. *Nat Med* 2001;7:687–92
94. Sata M, Nishimatsu H, Osuga J, Tanaka K, Ishizaka N, Ishibashi S, Hirata Y, Nagai R. Statins augment collateral growth in response to ischemia but they do not promote cancer and atherosclerosis. *Hypertension* 2004;43:1214–20
95. Hakamada-Taguchi R, Uehara Y, Kuribayashi K, Numabe A, Saito K, Negoro H, Fujita T, Toyo-oka T, Kato T. Inhibition of hydroxymethylglutaryl-coenzyme A reductase reduces Th1 development and promotes Th2 development. *Circ Res* 2003;93:948–56
96. Wong WW, Dimitroulakos J, Minden MD, Penn LZ. HMG-CoA reductase inhibitors and the malignant cell: the statin family of drugs as triggers of tumor-specific apoptosis. *Leukemia* 2002;16:508–19